

Publications en lien avec les activités du GDR "Calculs sur réseau en Physique Subatomique"

References

- [1] C. Alexandrou *et al.* [European Twisted Mass Collaboration], "Light baryon masses with dynamical twisted mass fermions," *Phys. Rev. D* **78**, 014509 (2008) [arXiv:0803.3190 [hep-lat]].
- [2] R. Baron, S. Capitani, J. Carbonell, K. Jansen, Z. Liu, O. Pene and C. Urbach [ETM Collaboration], "Moments of meson distribution functions with dynamical twisted mass fermions," *PoS LAT2007*, 153 (2007) [arXiv:0710.1580 [hep-lat]].
- [3] C. Alexandrou *et al.* [ETM Collaboration], "Baryon masses with dynamical twisted mass fermions," *PoS LAT2007*, 087 (2007) [arXiv:0710.1173 [hep-lat]].
- [4] F. De Soto, J. Carbonell, C. Roiesnel, Ph. Boucaud, J. P. Leroy and O. Pene, "Two body scattering length of Yukawa model on a lattice," *Nucl. Phys. A* **790**, 410 (2007) [arXiv:hep-lat/0610086].
- [5] F. De Soto, J. Carbonell, C. Roiesnel, Ph. Boucaud, J. P. Leroy and O. Pene, "Yukawa model on a lattice: Two body states," *Eur. Phys. J. A* **31**, 777 (2007) [arXiv:hep-lat/0610084].
- [6] F. de Soto, J. Carbonell, C. Roiesnel, Ph. Boucaud, J. P. Leroy and O. Pene, "Nuclear models on a lattice," *Nucl. Phys. Proc. Suppl.* **164**, 252 (2007) [arXiv:hep-lat/0511009].
- [7] S. Dürr, Z. Fodor, J. Frison, C. Hoelbling, R. Hoffmann, S. D. Katz, S. Krieg, T. Kurth, L. Lellouch, T. Lippert, K.K. Szabo, G. Vulvert [BMW Collaboration] "Ab-initio Determination of the Light Hadron Spectrum", CPT-P032-2008
- [8] C. Bourrely, I. Caprini and L. Lellouch, "Model-independent description of $B \rightarrow \pi l \nu$ decays and a determination of $|V_{ub}|$," arXiv:0807.2722 [hep-ph].
- [9] G. .. Akemann, F. Basile and L. Lellouch, "Finite size scaling of meson propagators with isospin chemical potential," arXiv:0804.3809 [hep-lat].
- [10] S. Dürr, Z. Fodor, C. Hoelbling, R. Hoffmann, S. D. Katz, S. Krieg, T. Kurth, L. Lellouch, T. Lippert, K.K. Szabo, G. Vulvert [BMW Collaboration] "Scaling study of dynamical smeared-link clover fermions," arXiv:0802.2706 [hep-lat].
- [11] S. Dürr, Z. Fodor, C. Hoelbling, S. D. Katz, S. Krieg, T. Kurth, L. Lellouch, T. Lippert, K.K. Szabo, G. Vulvert [BMW Collaboration] "Chiral behavior of pseudo-Goldstone boson masses and decay constants in 2+1 flavor QCD," *PoS LAT2007*, 115 (2007) [arXiv:0710.4769 [hep-lat]].

- [12] S. Dürr, Z. Fodor, C. Hoelbling, S. D. Katz, S. Krieg, T. Kurth, L. Lellouch, T. Lipert, K.K. Szabo, G. Vulvert [BMW Collaboration] “Mixed action simulations: approaching physical quark masses,” PoS **LAT2007**, 113 (2007) [arXiv:0710.4866 [hep-lat]].
- [13] R. Babich, N. Garron, C. Hoelbling, J. Howard, L. Lellouch and C. Rebbi, “Diquark correlations in baryons on the lattice with overlap quarks,” Phys. Rev. D **76**, 074021 (2007) [arXiv:hep-lat/0701023].
- [14] R. Babich, N. Garron, C. Hoelbling, J. Howard, L. Lellouch and C. Rebbi, “Matrix elements and diquark correlations in quenched QCD with overlap fermions,” PoS **LAT2006**, 091 (2006) [arXiv:hep-lat/0610079].
- [15] R. Babich, N. Garron, C. Hoelbling, J. Howard, L. Lellouch and C. Rebbi, “ K^0 - \bar{K}^0 mixing beyond the standard model and CP-violating electroweak penguins in quenched QCD with exact chiral symmetry,” Phys. Rev. D **74**, 073009 (2006) [arXiv:hep-lat/0605016].
- [16] R. Babich *et al.*, “Light hadron and diquark spectroscopy in quenched QCD with overlap quarks on a large lattice,” JHEP **0601**, 086 (2006) [arXiv:hep-lat/0509027].
- [17] R. Babich *et al.*, “Light hadron spectroscopy in quenched QCD with overlap fermions,” PoS **LAT2005**, 043 (2006) [arXiv:hep-lat/0509182].
- [18] F. Belletti *et al.* [apeNEXT Collaboration], Nucl. Instrum. Meth. A **559**, 90 (2006).
- [19] Ph. Boucaud, J. P. Leroy, A. Le Yaouanc, J. Micheli, O. Pene and J. Rodriguez-Quintero, “On the IR behaviour of the Landau-gauge ghost propagator,” JHEP **0806**, 099 (2008) [arXiv:0803.2161 [hep-ph]].
- [20] Ph. Boucaud *et al.* [ETM collaboration], “Dynamical Twisted Mass Fermions with Light Quarks: Simulation and Analysis Details,” arXiv:0803.0224 [hep-lat].
- [21] Ph. Boucaud, J. P. Leroy, A. L. Yaouanc, J. Micheli, O. Pene and J. Rodriguez-Quintero, “IR finiteness of the ghost dressing function from numerical resolution of the ghost SD equation,” JHEP **0806**, 012 (2008) [arXiv:0801.2721 [hep-ph]].
- [22] P. Boucaud *et al.*, “Ghost and gluon leading power corrections in Landau gauge,” Nucl. Phys. Proc. Suppl. **164**, 221 (2007).
- [23] Ph. Boucaud *et al.* [ETM Collaboration], “Dynamical twisted mass fermions with light quarks,” Phys. Lett. B **650**, 304 (2007) [arXiv:hep-lat/0701012].
- [24] Ph. Boucaud *et al.*, “Divergent IR gluon propagator from Ward-Slavnov-Taylor identities?,” JHEP **0703**, 076 (2007) [arXiv:hep-ph/0702092].
- [25] Ph. Boucaud *et al.*, “Constraints on the IR behaviour of gluon and ghost propagator from Ward-Slavnov-Taylor identities,” Eur. Phys. J. A **31**, 750 (2007) [arXiv:hep-ph/0701114].

- [26] Ph. Boucaud *et al.*, “Is the QCD ghost dressing function finite at zero momentum?,” JHEP **0606**, 001 (2006) [arXiv:hep-ph/0604056].
- [27] D. Becirevic, Ph. Boucaud, V. Lubicz, G. Martinelli, F. Mescia, S. Simula and C. Tarantino, “Exploring twisted mass lattice QCD with the clover term,” Phys. Rev. D **74**, 034501 (2006) [arXiv:hep-lat/0605006].
- [28] Ph. Boucaud *et al.*, “Short comment about the lattice gluon propagator at vanishing momentum,” arXiv:hep-lat/0602006.
- [29] Ph. Boucaud *et al.*, “The infrared behaviour of the pure Yang-Mills Green functions,” arXiv:hep-ph/0507104.
- [30] Ph. Boucaud *et al.*, “Non-perturbative power corrections to ghost and gluon propagators,” JHEP **0601**, 037 (2006) [arXiv:hep-lat/0507005].
- [31] Ph. Boucaud *et al.*, “Asymptotic behavior of the ghost propagator in SU3 lattice gauge theory,” Phys. Rev. D **72**, 114503 (2005) [arXiv:hep-lat/0506031].
- [32] Ph. Boucaud, F. de Soto, J. P. Leroy, A. Le Yaouanc, J. Micheli, O. Pene and J. Rodriguez-Quintero, “Instanton traces in lattice gluon correlation functions,” arXiv:hep-ph/0505150.
- [33] A. Y. Lokhov *et al.*, “Non-perturbative approach to the Landau gauge gluodynamics,” arXiv:hep-lat/0601007.
- [34] D. Becirevic *et al.*, “Non-perturbatively renormalised light quark masses from a lattice simulation with $N(f) = 2$,” Nucl. Phys. B **734**, 138 (2006) [arXiv:hep-lat/0510014].
- [35] D. Becirevic *et al.*, “Non-perturbatively renormalized light quark masses with two dynamical fermions,” PoS **LAT2005**, 079 (2006) [arXiv:hep-lat/0509091].
- [36] D. Becirevic, B. Haas and F. Mescia, “Semileptonic D-decays and Lattice QCD,” PoS **LAT2007**, 355 (2007) [arXiv:0710.1741 [hep-lat]].
- [37] D. Becirevic, V. Lubicz and F. Mescia, “An estimate of the $B - \bar{c} K^*$ gamma form factor,” Nucl. Phys. B **769**, 31 (2007) [arXiv:hep-ph/0611295].
- [38] S. Nicolis, “Layered Phase Investigations,” PoS **LAT2007**, 273 (2007) [arXiv:0710.1714 [hep-lat]].
- [39] B. Blossier, “Lattice renormalisation of $O(a)$ improved heavy-light operators,” Phys. Rev. D **76**, 114513 (2007) [arXiv:0705.0283 [hep-lat]].
- [40] I. I. Bigi *et al.*, “Memorino on the ‘1/2 vs. 3/2 Puzzle’ in $\bar{B} \rightarrow l\bar{\nu} X_c$ – a Year Later and a Bit Wiser,” Eur. Phys. J. C **52**, 975 (2007) [arXiv:0708.1621 [hep-ph]].

- [41] B. Blossier *et al.* [European Twisted Mass Collaboration], “Light quark masses and pseudoscalar decay constants from Nf=2 Lattice QCD with twisted mass fermions,” *JHEP* **0804**, 020 (2008) [arXiv:0709.4574 [hep-lat]].
- [42] B. Blossier, G. von Hippel, T. Mendes, R. Sommer and M. Della Morte, “Efficient use of the Generalized Eigenvalue Problem,” arXiv:0808.1017 [hep-lat].